



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

The index case of SARS-CoV-2 in Scotland: a case report

Citation for published version:

Hill, DKJ, Russell, DCD, Clifford, DS, Templeton, DK, Mackintosh, DCL, Koch, DO & Sutherland, DRK
2020, 'The index case of SARS-CoV-2 in Scotland: a case report', *Journal of Infection*.
<https://doi.org/10.1016/j.jinf.2020.03.022>

Digital Object Identifier (DOI):

[10.1016/j.jinf.2020.03.022](https://doi.org/10.1016/j.jinf.2020.03.022)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of Infection

Publisher Rights Statement:

Author's peer reviewed manuscript as accepted for publication.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



The index case of SARS-CoV-2 in Scotland: a case reportDr Katherine J. Hill¹Dr Clark D. Russell^{1,2}Dr Sarah Clifford¹Dr Kate Templeton³Dr Claire L. Mackintosh¹Dr Oliver Koch¹Dr Rebecca K. Sutherland¹¹NHS Lothian, Regional Infectious Diseases Unit, Edinburgh, EH4 2XU.²University of Edinburgh Centre for Inflammation Research, The Queen's Medical Research Institute, Edinburgh EH16 4TJ³NHS Lothian, Diagnostic Virology Reference Laboratory, Royal Infirmary of Edinburgh, Edinburgh, EH16 4TJ

Corresponding author: Dr Katherine J Hill, k.hill@doctors.org.uk 0131 537 2871

Abstract

Since its identification in December 2019, SARS-CoV-2 has infected 125,048 persons globally with cases identified in 118 countries across all continents (1). We report on the Scottish index case of SARS-CoV-2 infection, the virus causing COVID-19.

Keywords

Coronavirus

Communicable diseases, emerging

Respiratory tract infections

Letter

We read with interest Lillie and colleagues' letter on the first patients with COVID-19 in the UK (1). Since its identification in December 2019, SARS-CoV-2 has infected 125,048 persons globally with cases identified in 118 countries across all continents (2). We report on the Scottish index case of SARS-CoV-2 infection, the virus causing COVID-19.

The patient, a 51-year-old male, contacted the Scottish telehealth service for advice on day +1 with a 24-hour history of fever and cough having returned to Scotland from northern Italy on day -2. His symptoms were in keeping with a "possible" case of COVID-19 according to the national case definition at the time. Community SARS-CoV-2 testing was arranged for day +2.

The patient had travelled to Italy on day -9 to watch a rugby match in Rome. He travelled with his partner and two friends in a private rental vehicle through Lombardy, Veneto and Tuscany, staying in private rental accommodation, before flying back to Scotland from Milan on day -2. He was not aware of any contact with cases of COVID-19.

On day 0 he developed fever, myalgia, malaise and sinusitis. This progressed to a cough productive of green sputum on day +1. His fever had subsided by day +2.

The patient remained in self isolation in the community whilst awaiting the results of SARS-CoV-2 PCR which was performed in the West of Scotland Specialist Virology Centre on a combined nose/throat swab. This sample tested positive for SARS-CoV-2 on day +3 with a threshold cycle (Ct) value of 36. An urgent teleconference was held between virology, public health, Scottish ambulance service and the receiving regional High Consequence Infectious Diseases (HCID) unit. The patient was subsequently transferred by Special Operations Response Team (SORT) ambulance to the HCID unit. The patient was escorted from the ambulance to a laminar flow room (with antechamber) on the first floor of the unit by medical staff in appropriate personal protective equipment (PPE). The patient was wearing a surgical face mask throughout transit. The ambulance staff returned to their base for doffing of PPE.

The patient was usually well and exercised regularly. He was a lifelong non-smoker. His medical history consisted only of hypertension. He had no known underlying lung disease. He took an ACE-inhibitor and self-medicated with paracetamol and non-steroidal anti-inflammatory drugs during his illness.

On admission, the patient was afebrile with a mild non-productive cough. He had no limitation in exercise tolerance during this acute illness and continued to exercise in his home whilst self-isolating. On assessment, lung fields were clear on auscultation and bilateral scleral injection was noted. Admission vital signs were: heart rate 79 beats per minute, blood pressure 171/105mmHg, respiratory rate 18, pulse oximetry 96% on air and temperature 36.5. The results of routine laboratory parameters are shown in Table 1. By day three of admission the patient had lymphocytopenia, which was found in 80.4% of patients with laboratory-confirmed non-severe COVID-19 in mainland China (3). Chest radiography was not clinically indicated. PCR testing for other respiratory viruses, *Mycoplasma pneumoniae* and *Chlamydia pneumoniae* on the throat swab was negative. Nose and throat samples for SARS-CoV-2 PCR were obtained daily to monitor viral shedding (Figure 1). SARS-CoV-2 PCR on urine, faeces and EDTA blood was negative. The patient was clinically well throughout the admission; no supportive care was required. He was discharged after 8 days, following two sequential nose and throat swabs negative for SARS-CoV-2 by PCR. The initial viral sample was sequenced by the MRC-University of Glasgow Centre for Virus Research, mapped against the Wuhan-Hu-1 reference genome and aligned with a collection of global SARS-CoV-2 sequences, with the resulting analysis released open access. Phylogenetic analysis demonstrated that Scotland/CVR01/2020 belonged to a clade of isolates all acquired in northern Italy (4).

Public health undertook contact tracing and testing and no further cases were identified

A large case series from China has reported that the majority of COVID-19 cases (81%) had mild disease only, with risk of severe disease and mortality higher in older patients with pre-existing medical conditions (5). The patient described here also had a mild and self-limiting illness. Although asymptomatic and mild manifestations of previous epidemic-associated coronaviruses (SARS and MERS) occur (6,7), these appear to be substantially more common in COVID-19. This poses a challenge for the early identification and isolation of cases, to limit transmission. In the “contain” phase of the UK COVID-19 response, early case identification was reliant on epidemiologic (not severity) criteria. The public health guidance at the time of this case was that travellers to Veneto and Lombardy were considered at risk. The patient had travelled to these areas but in private transport, staying in the area briefly and having limited interactions with residents. Therefore, whilst fulfilling ‘at risk’ criteria, from a practical perspective the risk seemed low.

The kinetics of SARS-CoV-2 viral load have not yet been well characterised in large numbers of patients. Prolonged nasopharyngeal viral shedding was reported in a series of 18 patients from Singapore, with a median of 12 days between first and last positive samples and an initial large decline in viral load followed by a slower decay of residual low-level virus (8). Similar kinetics were observed in 17 patients from China, where results were stratified by site tested (nose/throat) suggesting quantitatively greater and more prolonged shedding from the nose (9). The kinetics of shedding in this case match this pattern, with more prolonged nasal shedding (Figure).


National public health guidance on PPE requirements for managing cases of COVID-19 in infectious disease units stipulates use of disposable gowns, two pairs of disposable gloves, visor and FFP3 respirator (10). These guidelines currently do not recommend use of specific footwear, so it is concerning that SARS-CoV-2 was detected by PCR from the shoe front of a healthcare worker caring for COVID-19 patients (11). For this case we used disposable boot covers over plastic clogs.

As the SARS-CoV-2 outbreak continues, we are learning more about the clinical manifestations of the infection and the logistics of containing and managing cases. This case highlights the need to have a low index of suspicion for diagnosis of COVID-19 and for early isolation.

Table 1: Routine laboratory parameters

Parameter	Normal	Day 1	Day 2	Day 3
Total white cell count ($\times 10^9/L$)	4–11	4.9	4.5	4.6
Neutrophil count ($\times 10^9/L$)	2–7.5	2.5	2.1	2.4
Lymphocyte count ($\times 10^9/L$)	1.5–4	1.6	1.8	1.3
Monocyte count ($\times 10^9/L$)	0.2–0.8	0.84	0.58	0.82
Platelet count ($\times 10^9/L$)	150–400	239	242	240
C-reactive protein (mg/L)	0–5	4	2	3

Renal and liver biochemistry values were normal.

Figure 1: SARS-CoV-2 PCR threshold cycle values during hospital admission

Note on day 2 a combined throat and nose swab was performed. Negative PCR is plotted as Ct value of 40 (dotted line).

Contributors

KH, CDR, SC, RS conceived of the correspondence. KH and CDR collected the data. KT contributed to the section on virological testing. All authors contributed to the writing of the final version of the article.

Declaration of Competing Interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We are grateful to the patient for providing informed consent to publish this report. Our thanks go to nursing staff, laboratory and medical colleagues in NHS Lothian who contributed directly or indirectly to patient care. We are grateful to colleagues in public health who contribute their expertise to management of these cases. This case report did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. There was no writing assistance.

References

- (1) Lillie PJ, Samson A, Li A, Adams K, Capstick R, Barlow GD, et al. Novel coronavirus disease (Covid-19): the first two patients in the UK with person to person transmission. *J Infect*. Published online ahead of print Feb 28 2020. doi: 10.1016/j.jinf.2020.02.020.
- (2) WHO. Coronavirus diseases 2019 (COVID-19): Situation report - 52. Available from: https://www.who.int/docs/default-source/coronaviruse/20200312-sitrep-52-covid-19.pdf?sfvrsn=e2bfc9c0_2 [Accessed March 14 2020].
- (3) Guan W, Zheng-yi N, Hu Y, Liang W, Ou C, He J, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med*. Published online February 28, 2020. doi: 10.1056/NEJMoa2002032.
- (4) Robertson DL. First report of COVID-19 in Scotland. Available from: <http://virological.org/t/first-report-of-covid-19-in-scotland/412> [Accessed March 13 2020].
- (5) Wu Z, McGoogan JM. Characteristics of and Important Lessons from the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72 314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA*. Published online February 24 2020. DOI:10.1001/jama.2020.2648
- (6) Wilder-Smith A, Telesman MD, Heng BH, Earnest A, Ling AE, Leo YS. Asymptomatic SARS coronavirus infection among healthcare workers, Singapore. *Emerg Infect Dis*. 2005 Jul;11(7):1142-1145. DOI: 10.3201/eid1107.041165.
- (7) Arabi YM, Balkhy HH, Hayden FG, Bouchama A, Luke T, Baillie JK, et al. Middle East Respiratory Syndrome. *N Engl J Med*. 2017 Feb 9;376(6):584-594. doi: 10.1056/NEJMSr1408795.
- (8) Young BE, Ong SWX, Kalimuddin S, Low JG, Tan SY, Loh J, et al. Epidemiologic Features and Clinical Course of Patients Infected With SARS-CoV-2 in Singapore. *JAMA*. Published online Mar 3 2020. doi: 10.1001/jama.2020.3204
- (9) Zou L, Ruan F, Huang M, Liang L, Huang H, Hong Z, et al. SARS-CoV-2 Viral Load in Upper Respiratory Specimens of Infected Patients. *N Engl J Med*. Published online Feb 19 2020. doi: 10.1056/NEJMc2001737
- (10) Health Protection Scotland. Infection prevention and control advice for acute care settings: Acute respiratory illness from novel or emerging pathogens (Coronavirus (COVID-19)) [Version 9.0]. Published March 10 2020. Available from: https://hpspubsrepo.blob.core.windows.net/hps-website/nss/2526/documents/1_infection-control-acute-care.pdf [Accessed 13 March 2020]
- (11) Ong SWX, Tan YK, Chia PY, Lee TH, Ng OT, Wong MSY, et al. Air, surface environmental, and personal protective equipment contamination by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) from a symptomatic patient. *JAMA*. Published online Mar 4 2020. doi: 10.1001/jama.2020.3227